



2010 PQRI Measures – Infectious Disease

*CINA Rec = Measures that are the simplest to pull data on; CINA recommends these measures for your specialty

**Optional: Measures that CINA can report on if there is adequate data in the EMR

Measure	Title	Description	*CINA Rec.	**Optional
83	Hepatitis C: Testing for Chronic Hepatitis C – Confirmation of Hepatitis C Viremia	Percentage of patients aged 18 years and older with a diagnosis of hepatitis C seen for an initial evaluation who had HCV RNA testing ordered or previously performed		✓
86	Hepatitis C: Antiviral Treatment Prescribed	Percentage of patients aged 18 years and older with a diagnosis of chronic hepatitis C who were prescribed peginterferon and ribavirin therapy within the 12-month reporting period		✓
110	Preventive Care and Screening: Influenza Immunization for Patients 50 Years and older	Percentage of patients aged 50 years and older who received an influenza immunization during the flu season (September through February)	✓	
114	Preventive Care and Screening: Inquiry Regarding Tobacco Use	Percentage of patients aged 18 years or older who were queried about tobacco use one or more times within 24 months	✓	
115	Preventive Care and Screening: Advising Smokers and Tobacco Users to Quit	Percentage of patients aged 18 years and older and are smokers or tobacco users who received advice to quit smoking		✓
124	Health Information Technology (HIT): Adoption/Use of Electronic Health Records (EHR)	Documents whether provider has adopted and is using health information technology. To qualify, the provider must have adopted and be using a certified/qualified EHR	✓	
159	HIV/AIDS: CD4+ Cell Count or CD4+ Percentage	Percentage of patients aged 6 months and older with a diagnosis of HIV/AIDS for whom a CD4+ cell count or CD4+ cell percentage was performed at least once every 6 months	✓	
161	HIV/AIDS: Adolescent and Adult Patients with HIV/AIDS Who Are Prescribed Potent Antiretroviral Therapy	Percentage of patients with a diagnosis of HIV/AIDS aged 13 years and older: who have a history of a nadir CD4+ cell count below 350/mm ³ or who have a history of an AIDS-defining condition, regardless of CD4+ cell count; or who are pregnant, regardless of CD4+ cell count or age, who were prescribed potent antiretroviral therapy	✓	
205	HIV/AIDS: Sexually Transmitted Disease Screening for Chlamydia and Gonorrhea	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS for whom chlamydia and gonorrhea screenings were performed at least once since the diagnosis of HIV infection	✓	
206	HIV/AIDS: Screening for High Risk Sexual Behaviors	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS who were screened for high risk sexual behaviors at least once within 12 months		✓
207	HIV/AIDS: Screening for Injection Drug Use	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS who were screened for injection drug use at least once within 12 months		✓
208	HIV/AIDS: Sexually Transmitted Disease Screening for Syphilis	Percentage of patients aged 13 years and older with a diagnosis of HIV/AIDS who were screened for syphilis at least once within 12 months	✓	