



2010 PQRI Measures – Urology

*CINA Rec = Measures that are the simplest to pull data on; CINA recommends these measures for your specialty

**Optional: Measures that CINA can report on if there is adequate data in the EMR

Measure	Title	Description	*CINA Rec.	**Optional
48	Urinary Incontinence: Assessment of Presence or Absence of Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older who were assessed for the presence or absence of urinary incontinence within 12 months		✓
49	Urinary Incontinence: Characterization of Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence whose urinary incontinence was characterized at least once within 12 months		✓
50	Urinary Incontinence: Plan of Care for Urinary Incontinence in Women Aged 65 Years and Older	Percentage of female patients aged 65 years and older with a diagnosis of urinary incontinence with a documented plan of care for urinary incontinence at least once within 12 months	✓	
102	Prostate Cancer: Avoidance of Overuse of Bone Scan for Staging Low-Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at low risk of recurrence receiving interstitial prostate brachytherapy, OR external beam radiotherapy to the prostate, OR radical prostatectomy, OR cryotherapy who did not have a bone scan performed at any time since diagnosis of prostate cancer		✓
104	Prostate Cancer: Adjuvant Hormonal Therapy for High-Risk Prostate Cancer Patients	Percentage of patients, regardless of age, with a diagnosis of prostate cancer at high risk of recurrence receiving external beam radiotherapy to the prostate who were prescribed adjuvant hormonal therapy (GnRH agonist or antagonist)		✓
105	Prostate Cancer: Three-Dimensional (3D) Radiotherapy	Percentage of patients with a diagnosis of clinically localized prostate cancer receiving external beam radiotherapy as a primary therapy to the prostate with or without nodal irradiation (no metastases; no salvage therapy) who receive three-dimensional conformal radiotherapy (3D-CRT) or intensity modulated radiation therapy (IMRT)		✓
114	Preventive Care and Screening: Inquiry Regarding Tobacco Use	Percentage of patients aged 18 years or older who were queried about tobacco use one or more times within 24 months	✓	
115	Preventive Care and Screening: Advising Smokers and Tobacco Users to Quit	Percentage of patients aged 18 years and older and are smokers or tobacco users who received advice to quit smoking		✓
124	Health Information Technology (HIT): Adoption/Use of Electronic Health Records (EHR)	Documents whether provider has adopted and is using health information technology. To qualify, the provider must have adopted and be using a certified/qualified EHR	✓	